DENTAL PROGRAMS CONSENT FOR ENDODONTIC (ROOT CANAL) SERVICES

		Date of Birth	
I hereby authorize Dr,		, and any other dentists of	to perform an
endodontic (root canal) procedure on tooth (teeth) #, and I understand that			
this is an \square elective / \square urgent / \square emergency procedure (check one).			
become tooth th	nal therapy is indicated when the pulp chamber of a infected. The procedure is accomplished when the at will allow it to be disinfected and then sealed with a subsequent passage of bacteria into or out of the to	e dentist creates a small opening th an inert rubber-like substance	in the biting surface of the
increase	been informed that the risks to my health if this proceed pain, swelling, loss of the tooth (teeth), loss of ot n, cyst formation, and/or deterioration of general health.	her teeth nearby, loss of the sup	
certain	been informed of possible alternative methods of tre inherent and potential risks in any treatment or proc the following:		
•	A failure to completely eliminate the infection requiring retreatment, root surgery or removal of the tooth at a later date;		
•	Post-operative pain, swelling, bruising, and/or limited jaw opening that may persist for several days; Separation (breakage) of an instrument within the canal during treatment. Broken instrument tips are typically allowed to remain in the canal, and only rarely are they the cause of subsequent problems. If removal is indicated the patient may be referred to an endodontic specialist.		
•	Perforation of the root from within the canal can occur requiring additional treatment by a specialist. Such complications will occasionally result in the loss of the tooth. Damage to nerves supplying the teeth resulting in temporary or, in rare instances, permanent numbness or		
•	tingling of the lip, chin, or other areas of the jaws or face: Inability to adequately clean the canal(s) due to unforeseen calcified obstructions or severely bent roots. Under certain circumstances the patient may be referred to a specialist for successful completion of the procedure. Loss of the tooth may occur:		
•	A fracture of the treated tooth, occurring during or after endodontic treatment. Treated teeth sometimes break due to the tooth's loss of strength resulting from the procedure. In most cases a crown is recommended after treatment to prevent such an occurrence.		
require	eatment has begun, it is essential that it be complete from 1-5 appointments. Also, I understand that sucreated tooth.		
	stand the recommended treatment, the risks of such and the consequences of doing nothing.	treatment, alternative treatment	s should any
Patient's Signature		Date	
Parent or Legal Guardian Signature		Date	
Witness or Interpreter		Date	
Dentist's Signature		Date	